arXiv:1104.3650v1 [math-ph] 19 Apr 2011

Unified Analytic Electron Interaction Integrals Over Slater Orbitals for Diatomic Molecules

E.V.Rothstein

evrothstein@gmail.com

A unified formula for the analytical evaluation of two-center exchange, hybrid and coulombic type integrals is presented.

INTRODUCTION

A formula for the analytic evaluation of two-center electron interaction integrals is presented. The terms in the sum have been checked[1]

FORMULA

$$\Phi_{ic}(j) = (-)^{(m_i-|m_i|)/2}(2\delta_i)^{n_i+\frac{1}{2}}\left[\frac{(2l_i+1)(l_i-|m_i|)!}{4\pi(2n_i)!(l_i+|m_i|)!}\right]^{\frac{1}{2}} \times e^{im_i\phi_j} r_{jc}^{n_i-1} e^{-\delta_i r_{jc}} P_{l_i}^{|m_i|}(cos\theta_{jc})$$

i = orbital number with quantum numbers n_i, l_i, m_i and screening constant δ_i . c = center, centered on nucleus a or b . $\Phi_{ic}(j)$ = Normalized Slater Orbital for electron j, in spherical coordinates centered on center c.

I = integral to be evaluated known as the two-center exchange integral

$$I = \langle \Phi_{1a}(1)\Phi_{3b}(1)\frac{1}{r_{12}}\Phi_{2a}(2)\Phi_{4b}(2)\rangle$$

$$I = \delta(\sum m_i)\delta(|\sigma| - |m_2+m_4|)W\sum_k C_k(g_{1k}, g_{2k}, r_{1k}, r_{2k}) \times \sum_{\mu=|\sigma|}(2\mu+1)B(\mu, g_{1k}, \beta_1, |\sigma|)B(\mu, g_{2k}, \beta_2, |\sigma|)A(\mu, r_{1k}, r_{2k}, \alpha_1, \alpha_2, |\sigma|) \tag{1}$$

The δ in [Eq. (1)] is the Kronecker delta. The first sum is finite over k (the set of $g_{1k}, g_{2K}, r_{1k}, r_{2k}$ generated by the four orbitals) . The second sum is finite if either or both the β 's = 0 . If $\beta_1 = 0$ the sum over μ is restricted so the $\mu + |\sigma| + g_{1k}$ is even and the upper limit of the summation is $g_{1k} + |\sigma|$. If neither of the β's = 0 the upper limit is infinite. Some notation: $P_l^{|m|}(cos\theta)$ = Associated Legendre function of the first kind. (see Ref. 2 , 8.6.6) R = internuclear distance between a and b nuclei.

$$\begin{aligned}
&\alpha_1 = \frac{R}{2}(\delta_1 + \delta_3); \quad \alpha_2 = \frac{R}{2}(\delta_2 + \delta_4); \quad \beta_1 = \frac{R}{2}(\delta_1 - \delta_3); \\
&\quad \beta_2 = \frac{R}{2}(\delta_2 - \delta_4); \quad W = R^{\sum n_i + 1} \Pi \delta_i^{n_i + 1/2} \\
&W = \delta_1 (\alpha_1 + \beta_1)^{n_1 - 1/2} (\alpha_1 - \beta_1)^{n_3 + 1/2} (\alpha_2 + \beta_2)^{n_2 + 1/2} (\alpha_2 - \beta_2)^{n_4 + 1/2} \\
&W = \frac{1}{R} (\alpha_1 + \beta_1)^{n_1 + 1/2} (\alpha_1 - \beta_1)^{n_3 + 1/2} (\alpha_2 + \beta_2)^{n_2 + 1/2} (\alpha_2 - \beta_2)^{n_4 + 1/2}
\end{aligned}$$

$$\begin{aligned}
&\sum_k C_k(g_{1k}, g_{2k}.r_{1k}, r_{2k}) = (-)^{l_3 - |m_3| + l_4 - |m_4|} \left(\frac{1}{2}\right) \sum_{a_1=0}^{n_1 - l_1} \sum_{b_1=0}^{n_3 - l_3} \\
&\times \sum_{a_2=0}^{n_2 - l_2} \sum_{b_2=0}^{n_4 - l_4} (-)^{b_1 + b_2} \binom{n_1 - l_1}{a_1} \binom{n_3 - l_3}{b_1} \binom{n_2 - l_2}{a_2} \binom{n_4 - l_4}{b_2} \\
&\times \prod_{i=1}^{4} \left\{ \left[\frac{(2l_i + 1)(l_i - |m_i|)!(l_i + |m_i|)!}{(2n_i)!} \right]^{1/2} \sum_{s_i=0}^{[(l_i - |m_i|)/2]} \sum_{p_i=0}^{2s_i} \right. \\
&\times \sum_{q_i=0}^{l_i - |m_i| - 2s_i} \frac{(-)^{s_i + q_3 + q_4 + p_3 + p_4}}{2^{l_i} l_i!} \binom{2l_i - 2s_i}{l_i - |m_i| - 2s_i} \binom{l_i}{s_i} \\
&\times \binom{l_i - |m_i| - 2s_i}{q_i} \binom{2s_i}{p_i} \} \sum_{c_1=0}^{(|m_1| + |m_3| - |\sigma|)/2} \sum_{c_2=0}^{(|m_2| + |m_4| - |\sigma|)/2} \\
&\times \sum_{d_i=0}^{(|m_1| + |m_3| - |\sigma|)/2} \sum_{d_2=0}^{(|m_2| + |m_4| - |\sigma|)/2} (-)^{c_1 + c_2 + d_1 + d_2} \\
&\times \binom{(|m_1| + |m_3| - |\sigma|)/2}{c_1} \binom{(|m_1| + |m_3| - |\sigma|)/2}{d_1} \binom{(|m_2| + |m_4| - |\sigma|)/2}{c_2} \\
&\qquad\qquad \times \binom{(|m_2| + |m_4| - |\sigma|)/2}{d_2}. \quad (2)
\end{aligned}$$

The square brackets in the upper limit of summation over s_i is $\frac{1}{2}(l_i - |m_i|)$ or $\frac{1}{2}(l_i - |m_i| - 1)$ whichever is an integral.

$$g_1 = p_1 + p_3 + q_1 + q_3 + a_1 + b_1 + 2c_1,$$

$$g_2 = p_2 + p_4 + q_2 + q_4 + a_2 + b_2 + 2c_2,$$

$$r_1 = 2(s_1 + s_3 + d_1) + q_1 + q_3 - p_1 - p_3 + n_1 + n_3 - l_1 - l_3 - a_1 - b_1,$$

$$r_2 = 2(s_2 + s_4 + d_2) + q_2 + q_4 - p_2 - p_4 + n_2 + n_4 - l_2 - l_4 - a_2 - b_2, \text{ See footnote 3}$$

$$B(\mu, g, \beta, |\sigma|) = \left(\frac{-\partial}{\partial\beta}\right)^g \sqrt{2\pi}\, \frac{I_{\mu+1/2}(\beta)}{\beta^{|\sigma|+1/2}},$$

$I_{\mu+\frac{1}{2}}(\beta) =$ modified spherical Bessel function of the first kind

(see 10.2.2 Ref. 2)

$$\text{If } \beta = 0 \text{ then, } B(\mu, g, \beta, |\sigma|) = \frac{(-)^{\mu-|\sigma|} 2^{\mu+2} \binom{1+g+|\sigma|}{\frac{g+|\sigma|-\mu}{2}}}{(1+|\sigma|)! \binom{g+|\sigma|+1}{g} \binom{2+g+|\sigma|+\mu}{1+\frac{g+|\sigma|+\mu}{2}}}$$

$$\text{If } \beta \neq 0 \text{ then, } B(\mu, g, \beta, |\sigma|) = \frac{(-)^{\mu+1}}{\beta^{|\sigma|+1}} \sum_{k=0}^{\mu} \frac{\mu!}{(\mu-k)!} \binom{\mu+k}{k}$$

$$\times \frac{1}{(2\beta)^k} \sum_{j=0}^{g} \binom{g}{j} \binom{k+|\sigma|+j}{j} \frac{j!}{\beta^j} \left[(-)^{k+j+g+\mu+1} e^{\beta} + e^{-\beta}\right]$$

In practice this latter formula was not used. The ascending series formula proved to be more accurate (less roundoff errors) for $\beta \neq 0$ and $2k + \mu - |\sigma| \geq g$, then,

$$B(\mu, g, \beta, |\sigma|) = (-)^g 2^{\mu+1} \beta^{\mu-|\sigma|-g} \sum_{k=0}^{\infty} \frac{\beta^{2k} \binom{\mu+2k-|\sigma|}{g} \frac{g!}{k!\mu!(k+1)!}}{\binom{2\mu+2k+1}{\mu+k} \binom{\mu+k+1}{\mu}} \tag{3}$$

$$A(\mu, r_1, r_2, \alpha_1, \alpha_2, |\sigma|) = \left(\frac{-\partial}{\partial\alpha_1}\right)^{r_1} \left(\frac{-\partial}{\partial\alpha_2}\right)^{r_2} \int_1^\infty \int_1^\infty d\xi_1 d\xi_2$$

$$\times\, (\xi_1^2-1)^{|\sigma|/2}(\xi_2^2-1)^{|\sigma|/2} P_\mu^{|\sigma|}(\xi_{1<2}) Q_\mu^{|\sigma|}(\xi_{2>1}) e^{-\alpha_1\xi_1} e^{-\alpha_2\xi_2}$$

$$= \frac{e^{-(\alpha_1+\alpha_2)}}{2^{2\mu+1}} \left[\frac{(\mu+|\sigma|)!}{\mu!}\right]^2 \sum_{k=0}^{\left[\frac{\mu+|\sigma|}{2}\right]} (-)^k \binom{2\mu-2k}{\mu-|\sigma|} \binom{\mu}{k}$$

$$\times \sum_{p=0}^{\left[\frac{\mu+|\sigma|}{2}\right]} (-)^p \binom{2\mu-2p}{\mu-|\sigma|} \binom{\mu}{p} \{ \sum_{n_1=0}^{k_1} \frac{k_1!}{(k_1-n_1)!\,\alpha_1^{n_1+1}} \sum_{n_2=0}^{k_2} \frac{k_2!}{(k_2-n_2)!\,\alpha_2^{n_2+1}}$$

$$[\, \ln\left(\frac{2\alpha_1\alpha_2}{\alpha_1+\alpha_2}\right) + (-)^{n_1+n_2+k_1+k_2+1} e^{2(\alpha_1+\alpha_2)} E_1(2\alpha_1+2\alpha_2)$$

$$+ (-)^{k_1+n_1} e^{2\alpha_1} E_1(2\alpha_1) + (-)^{k_2+n_2} e^{2\alpha_2} E_1(2\alpha_2) + \gamma \,]$$

$$+ \sum_{n_2=1}^{k_2} \frac{1}{n_2} \sum_{j_2=0}^{k_2-n_2} \frac{k_2!}{(k_2-n_2-j_2)!} \sum_{n_1=0}^{k_1} [\, \frac{-k_1!}{(k_1-n_1)!} \frac{1}{\alpha_1^{n_1+1}\alpha_2^{n_2+j_2+1}}$$

$$+ \frac{k_1!}{(k_1-n_1)!} (-)^{k_2+n_2+j_2} \sum_{t=0}^{n_2-1} \frac{2^{n_2-t-1}}{(n_2-t-1)!\,\alpha_1^{n_1+1}\alpha_2^{j_2+t+2}}$$

$$+ \sum_{j_1=0}^{n_1} \frac{k_1!}{(n_1-j_1)!} (\, \binom{k_1+n_2-n_1-1}{k_1-n_1} \frac{1}{\alpha_2^{j_2+1}(\alpha_1+\alpha_2)^{k_1+n_2-n_1}\alpha_1^{j_1+1}}$$

$$+ \sum_{t=0}^{n_2-1} \frac{(-)^{k_2+n_2+j_2+1} 2^{n_2-t-1}}{(n_2-t-1)!} \binom{t+k_1-n_1}{t} \frac{1}{\alpha_2^{j_2+1}(\alpha_1+\alpha_2)^{k_1+t+1-n_1}\alpha_1^{j_1+1}} \,)\,]$$

$$+ \sum_{n_2=0}^{k_2} \frac{k_2!}{(k_2-n_2)!} \sum_{n_1=1}^{k_1} \frac{1}{n_1} \sum_{j_1=0}^{k_1-n_1} \frac{k_1!}{(k_1-n_1-j_1)!} [\, \frac{1}{\alpha_1^{j_1+1}\alpha_2^{n_2+1}(\alpha_1+\alpha_2)^{n_1}}$$

$$- \frac{1}{\alpha_1^{n_1+j_1+1}\alpha_2^{n_2+1}} + \sum_{t=0}^{n_1-1} \frac{(-)^{k_1+j_1+n_1} 2^{n_1-1-t}}{(n_1-t-1)!} (\, \frac{1}{\alpha_1^{j_1+t+2}\alpha_2^{n_2+1}}$$

$$+ \frac{(-)^{k_2+n_2+1}}{(\alpha_1+\alpha_2)^{t+1}\alpha_1^{j_1+1}\alpha_2^{n_2+1}} \,)\,]\,\} + \frac{e^{-(\alpha_1+\alpha_2)}(\mu+|\sigma|)!}{2^{2\mu}\mu!} \sum_{k=0}^{[\frac{\mu+|\sigma|}{2}]} (-)^k \binom{2\mu-2k}{\mu-|\sigma|}$$

$$\times \binom{\mu}{k} [\, \sum_{\kappa=1}^{|\sigma|} \frac{(-)^\kappa |\sigma|!}{\kappa} \binom{\mu+|\sigma|-\kappa}{\mu} \sum_{j=0}^{[\frac{\kappa-1}{2}]} \binom{\kappa}{\kappa-2j-1} \sum_{n=0}^{[\frac{\mu+|\sigma|-\kappa}{2}]} (-)^n$$

$$\times \binom{2\mu-2n}{\mu-|\sigma|+\kappa} \binom{\mu}{n} - \sum_{j=0}^{[\frac{\mu-|\sigma|-1}{2}]} \frac{(2\mu-4j-1)(\mu-2j-1+|\sigma|)!\,2^{2j+1}}{(2j+1)(\mu-j)(\mu-2j-1)!}$$

$$\times \sum_{n=0}^{[\frac{\mu+|\sigma|-2j-1}{2}]} (-)^n \binom{2(\mu-2j-1-n)}{\mu-2j-1-|\sigma|} \binom{\mu-2j-1}{n} \,]$$

$$\times\, [\, \sum_{n_1=0}^{k_1^o} \frac{k_1^o!}{(k_1^o - n_1)!} \sum_{n_2=0}^{f_2} \binom{n_1 + f_2 - n_2}{f_2 - n_2}$$

$$\times \sum_{j_2=0}^{n_2} \frac{f_2!}{(n_2 - j_2)!\, \alpha_2^{j_2+1} (\alpha_1 + \alpha_2)^{f_2+n_1-n_2+1}}$$

$$+ \sum_{n_2=0}^{k_2} \frac{k_2!}{(k_2 - n_2)!} \sum_{n_1=0}^{f_1} \binom{n_2 + f_1 - n_1}{f_1 - n_1}$$

$$\times \sum_{j_1=0}^{n_1} \frac{f_1!}{(n_1 - j_1)!\, \alpha_1^{j_1+1} (\alpha_1 + \alpha_2)^{f_1+n_2-n_1+1}}\,]\,.$$

$E_1(x) =$ Exponential integral (5.1.1. Ref. 2)

$\gamma =$ Euler's constant (table 1.1 Ref. 2)

$$k_1 = \mu + |\sigma| - 2p + r_1, \quad k_2 = \mu + |\sigma| - 2k + r_2,$$

$$k_1^o = \mu + |\sigma| - 2k + r_1, \quad f_1 = \mu + |\sigma| - 2(n + j) - 1 + r_1,$$

$$f_2 = \mu + |\sigma| - 2(n + j) - 1 + r_2, \quad (4)$$

1. E.R.Chan, Chem. Phys. Lett., 23 ,99 (1973) .
2. M. Abramowitz and I.A.Stegun, editors, *Handbook of Mathematical Functions*, (Dover, New York)
3.

$$HybridIntegral = \langle \Phi_{1a}(1)\Phi_{3a}(1)\frac{1}{r_{12}}\Phi_{2a}(2)\Phi_{4b}(2)\rangle$$

$$\alpha_1 = \beta_1 = \frac{R}{2}(\delta_1 + \delta_3);\ \alpha_2 = \frac{R}{2}(\delta_2 + \delta_4);\ \beta_2 = \frac{R}{2}(\delta_2 - \delta_4);$$

$$CoulombIntegral = \langle \Phi_{1a}(1)\Phi_{3a}(1)\frac{1}{r_{12}}\Phi_{2b}(2)\Phi_{4b}(2)\rangle$$

$$\alpha_1 = \beta_1 = \frac{R}{2}(\delta_1 + \delta_3);\ \alpha_2 = -\beta_2 = \frac{R}{2}(\delta_2 + \delta_4);$$

These changes result in changes in the set of $g_{1k}, g_{2k}, r_{1k}.r_{2k}.C_k$.